

חצין

<p>PATIENT PACKAGE INSERT IN ACCORDANCE WITH THE PHARMACISTS' REGULATIONS (PREPARATIONS) - 1966</p> <p>Read this package insert carefully in its entirety before using this medicine. The format of this leaflet was determined by the Ministry of Health and its content was checked and approved in January 2006</p> <p>CORSODYL™ MINT MOUTHWASH</p> <p>Composition: Chlorhexidine Gluconate 0.2% w/v.</p> <p>Inactive ingredients: Sorbitol Sol., Ethanol, Polyoxyethylated Hydrogenated Castor Oil, Peppermint Oil, Purified Water.</p> <p>Therapeutic group: Antiseptics.</p> <p>Therapeutic activity: Antibacterial solution for the disinfection of the mouth. Influential on the inhibition of gingivitis. To be used after surgery of the gums and in the treatment of mouth infections and ulcerations. This solution does not prevent the formation of dental plaque and dental calculus (tarar). That is why it is important to clean teeth thoroughly by correct brushing and by using dental floss.</p> <p>When should the preparation not be used?</p> <p>[Do not use this medicine if you are sensitive to any of its ingredients.]</p> <p>Do not use this medicine without consulting a physician before starting treatment:</p> <p>If you are pregnant or if you are breastfeeding.</p> <p>Warnings: If you are sensitive to any type of food or medicine or to chlorhexidine or to any disinfecting agent containing chlorhexidine, inform your physician immediately.</p> <p>Attention: Do not swallow! Do not dilute with water. Avoid contact of this preparation with eyes and ears.</p> <p>Storage: In a cool place. Even if kept in their original container and stored as recommended, medicines may be kept for a limited period only. Please note the expiry date of the medicine! In case of doubt, consult the pharmacist who dispensed the medicine to you. Do not store different medications in the same package.</p> <p>License number: 029 96 21599 D0</p> <p>Manufacturing site: SmithKline Beecham PLC, UK.</p> <p>License Holder and Importer: GlaxoSmithKline (Israel) Ltd., 25 Hazel Street, Petach-Tikva.</p>	<p>assistant.</p> <p>Transient changes in taste can occur. Corsodyl™ has a bitter taste, but do not rinse your mouth with water after using Corsodyl™ as it will increase the bitterness and reduce the effectiveness of the medication.</p> <p>Side effects that require special attention: Local irritation - (rare), stop treatment and refer to your physician. In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your physician immediately.</p> <p>Do not induce vomiting unless explicitly instructed to do so by a physician!</p> <p>Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.</p> <p>Attention: Do not swallow! Do not dilute with water. Avoid contact of this preparation with eyes and ears.</p> <p>Directions for use: Fill the bottle cap with Corsodyl™ solution up to the marked line (10 ml) and rinse the mouth twice a day for half a minute, immediately after brushing your teeth. To assist in the treatment of gingivitis: It is recommended to use Corsodyl™ for a month. For the treatment of mouth ulcers: Corsodyl™ should be used for an additional 2 days after the area has healed.</p> <p>After gum surgery or an extraction: Rinse with Corsodyl™ according to the doctor's instructions. Do not exceed the recommended dose.</p>	<p>Avoid poisoning! This medicine, and all other medicines, must be stored in a safe place out of the reach of children and/or infants, to avoid poisoning. If you have taken an overdose, or if a child has accidentally swallowed the medicine, proceed immediately to a hospital emergency room and bring the package of the medicine with you.</p> <p>Side effects that require special attention: Local irritation - (rare), stop treatment and refer to your physician. In the event that you experience side effects not mentioned in this leaflet, or if there is a change in your general health, consult your physician immediately.</p> <p>Do not induce vomiting unless explicitly instructed to do so by a physician!</p> <p>Do not take medicines in the dark! Check the label and the dose each time you take your medicine. Wear glasses if you need them.</p> <p>Attention: Do not swallow! Do not dilute with water. Avoid contact of this preparation with eyes and ears.</p> <p>Directions for use: Fill the bottle cap with Corsodyl™ solution up to the marked line (10 ml) and rinse the mouth twice a day for half a minute, immediately after brushing your teeth. To assist in the treatment of gingivitis: It is recommended to use Corsodyl™ for a month. For the treatment of mouth ulcers: Corsodyl™ should be used for an additional 2 days after the area has healed.</p> <p>After gum surgery or an extraction: Rinse with Corsodyl™ according to the doctor's instructions. Do not exceed the recommended dose.</p>	<p>ИСКРУЧКАЯ ПОРИНКЕНИЮ В СООТВЕТСТВИИ С ФАРМАЦЕТИЧЕСКИМИ СТАНДАРТАМИ (ПРЕПАРАТЫ) - 1966</p> <p>Внимательно и полностью прочтите инструкцию перед приемом препарата</p> <p>Формат настоящей инструкции установлен Министерством здравоохранения, содержание проверено и утверждено в январе 2006</p> <p>КОРСОДИЛ™ ОПОЛАСКИВАТЕЛЬ ДЛЯ ПОЛОСТИ РТА</p> <p>Состав: Хлоргексидина глюконат 0.2% w/v.</p> <p>Вспомогательные вещества: Sorbitol Sol., Ethanol, Polyoxyethylated Hydrogenated Castor Oil, Peppermint Oil, Purified Water.</p> <p>Побочные эффекты: Во время курса лечения, наряду с желаемым действием препарата, могут возникнуть побочные эффекты, такие как легкое ощущение горячего в ротовой полости. Препарат также может вызвать повышенное отложение зубного камня. Чистые зубы и чистая зубная паста перед применением препарата, можно обслабить эти явления. Если же все-таки произошло ощущение, оно может быть удалено чисткой зубов специалистом по гигиене полости рта.</p>	<p>налета и зубного камня, поэтому важно тщательно и правильно чистить зубы и пользоваться зубной нитью.</p> <p>Противопоказания к применению:</p> <p>[Противопоказано применение препарата при повышенном чувствительности к одному из компонентов препарата.]</p> <p>Не применяйте препарат без предварительной консультации врача: В период беременности или коррекции грудью.</p> <p>Предостережения: До начала приема препарата следует сообщить врачу о повышенной чувствительности к каким-либо продуктам питания или лекарственным препаратам, или хлоргексидину или к дезинфицирующему средству, содержащему хлоргексидин.</p> <p>Внимание: Не глотать! Не разбавлять водой. Избегайте контакта препарата с глазами и ушами.</p> <p>Указания к применению: Наполните крышку флакона раствором Корсодил до линии отметки (10 мл).</p> <p>Полоскайте полость рта в течение полинуты дневки в день сразу после чистки зубов.</p> <p>В качестве вспомогательного средства при воспалении десен: рекомендуется применять Корсодил в течение месяца.</p> <p>Для лечения язвочек в полости рта: рекомендуется применять Корсодил в течение двух дополнительных дней после захвивания.</p> <p>После операций на деснах или хирургического</p>	<p>esk</p> <p>לשיטיפת פה טכניון אוניברסיטת תל אביב הפקולטה לרפואה dept. of Dentistry DENTAL HYGIENE DEPARTMENT</p> <p>700 ml</p>
---	--	---	---	---	--

גנ